Immunization

NO FLU SHOT

Definition: Respondents ages 65 and older who have not had an influenza vaccination within the past 12 months.

Prevalence of No Flu Shot

- South Dakota 25.9%
- o Nationwide median 30.4%


Healthy People 2010 Objective

Decrease the proportion of adults age 65 years and older who are not vaccinated annually against influenza to 10 percent.

Trend Analysis

Overall, the percent of respondents aged 65 and older who have not had an annual flu shot has been decreasing. However, since 2003 it has been increasing. South Dakota has not yet met the *Healthy People 2010 Objective* of 10 percent.

Figure 50
Percent of Respondents, Ages 65 and Older, Who Have Not Had a Flu Shot
Within the Past 12 Months, 1993-1995, 1997, and 1999-2006



Demographics

Gender There is no significant gender difference shown in the lack of a flu shot for those

65 and older or for any demographic group between the two.

Age There is a significant decrease in the percentage of those with no flu shot from the

65-74 age group to the 75+ age group. This decrease is more evident in males.

Race There is insufficient data to make any racial comparisons.

Region There are no significant regional differences exhibited by the available data.

Household The lack of a flu shot seems to generally decrease as household income increases. **Income**

Education The lack of a flu shot seems to generally decrease as education levels increase.

Employment Those who are employed for wages or self-employed demonstrate a very high prevalence of no flu shot, while those who are a homemaker or retired show a

very low prevalence.

Marital Those who are divorced or separated exhibit a very high prevalence of no flu **Status** shot, while those who are married or widowed show a very low prevalence.

Table 62 Respondents, Ages 65 and Older, Who Have Not Had a Flu Shot Within the Past 12 Months, 2006 Total Male Female # Resp. # Resp. % 95% CI # Resp. % 95% CI % 95% CI 25.9 1,283 24.7 **Total** 1.950 (23.7-28.2)27.6 (23.9-31.5)(22.1-27.4)667 <u>Age</u> 18-24 25-34 35-44 --_ 45-54 55-64 30.0 (26.8-33.5) 35.1 (29.7-40.9) 578 25.7 (21.9-29.9) 941 363 65-74 (14.9-24.6)23.9 75 +1,009 22.1 (19.3-25.1)304 19.3 705 (20.5-27.7)Race (24.0-28.6) (24.1-31.9) 1,208 (22.3-27.8)1,835 26.2 627 27.8 25.0 White American Indian Region Southeast 134 417 23.8 (19.8-28.4)24.9 (18.1-33.2)283 23.1 (18.4-28.6)542 24.3 (20.5-28.5)178 29.7 (23.0-37.3)364 20.5 (16.4-25.4)Northeast $(\overline{18.0-32.3})$ (22.2-30.8)(22.6-33.3) 446 26.3 148 24.4 298 27.6 Central 30.5 251 30.4 399 (25.8-35.6) 148 30.6 (23.4-39.0)(24.7-36.7) West American Indian Counties 31.9 (23.9-41.2)146 **Household Income** Less than \$10,000 120 29.7 (20.7-40.6)\$10,000-\$14,999 205 29.1 (22.5-36.6)\$15,000-\$19,999 212 27.8 (21.3-35.3)* \$20,000-\$24,999 235 25.4 (19.5-32.4)119 (19.2-36.9)\$25,000-\$34,999 289 28.8 (23.2-35.2)27.1 170 30.4 (23.0-39.1)\$35,000-\$49,999 240 26.5 (20.7-33.4)123 29.1 (20.9-38.9)117 23.0 (15.7-32.4)\$50,000-\$74,999 122 21.4 (14.7-30.2)* * * \$75,000+ **Education** 223 29.5 100 28.0 123 (22.3-41.6) (23.0-36.9)(19.2-38.8)31.2 8th Grade or Less 29.3 158 (21.6-38.5)Some High School 242 31.5 (25.3-38.4) 512 21.4 25.4 (17.7-25.5)High School or G.E.D. 754 (22.1-29.1)Some Post-High School 431 25.8 (21.3-30.9)105 24.8 (16.6-35.2)326 26.2 (21.1-32.1)380 23.6 (19.1-28.7) 167 21.9 (15.8-29.6)213 25.5 (19.5-32.7)College Graduate **Employment Status** * 210 35.7 (28.5-43.6)**Employed for Wages** (30.0-46.6)* * * * * Self-employed 178 38.0 * * Unemployed * * * * * * Homemaker 148 21.7 (15.3-29.9)* * * * * (20.5-25.6) 22.9 462 21.4 (17.5-25.8)880 24.1 (21.1-27.5)Retired 1.342 Unable to Work

Table 62 (continued) Respondents, Ages 65 and Older, Who Have Not Had a Flu Shot Within the Past 12 Months, 2006

	Total		Male			Female			
	# Resp.	%	95% CI	# Resp.	%	95% CI	# Resp.	%	95% CI
Marital Status									
Married/Unmarried Couple	889	25.1	(22.1-28.3)	445	27.3	(23.1-32.0)	444	22.3	(18.4-26.8)
Divorced/Separated	151	38.0	(29.0-47.9)	*	*	*	*	*	*
Widowed	843	25.3	(22.2-28.7)	124	21.8	(14.7-31.2)	719	26.0	(22.6-29.7)
Never Married	*	*	*	*	*	*	*	*	*


Note: *Results based on sample sizes less than 100 have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2006

National Statistics

The national median for respondents who have not had a flu shot within the past 12 months was 30.4 percent. South Dakota had 25.9 percent of respondents who have not had a flu shot within the past 12 months. Colorado had the lowest percent of respondents who have not had a flu shot within the past 12 months with 24.1 percent, while Nevada had the highest percent with 42.3 percent.

Figure 51
Nationally, Respondents Ages 65 and Older, Who Have Not Had a Flu Shot Within the
Past 12 Months 2006



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2006

Further Analysis

Following are data illustrating the percent of those, ages 65 and older, who have not had a flu shot for various health behaviors and conditions. For example, 42.1 percent of respondents who stated they currently smoke had no flu shot, while 23.5 percent of respondents who stated they were former smokers had no flu shot.

Table 63 No Flu Shot, Ages 65 and Older, Within the Past 12 Months for Selected Health Behaviors and Conditions, 2006

Health Behavior or Condition	# Respondents	% No Flu Shot	95% CI
Obese (BMI = $30.0+$)	409	24.3	19.9-29.4
Overweight (BMI = 25.0-29.9)	762	27.6	24.1-31.4
Recommended Weight (BMI = 18.5-24.9)	637	25.0	21.3-29.2
No Leisure Time Physical Activity	653	27.0	23.2-31.1
Leisure Time Physical Activity	1,292	25.5	22.9-28.3
Not Heard of "Healthy South Dakota" Program	1,089	26.6	23.7-29.7
Heard of "Healthy South Dakota" Program	762	24.0	20.7-27.6
Current Smoker	189	42.1	34.1-50.5
Former Smoker	712	23.5	201-27.2
Never Smoked	1,047	25.0	22.1-28.2
Smokeless Tobacco Use	*	*	*
No Smokeless Tobacco Use	1,863	25.4	23.2-27.7
Diabetes	294	19.6	15.0-25.3
No Diabetes	1,655	27.0	24.6-29.5
Current Asthma	134	23.3	16.2-32.3
Former Asthma	*	*	*
Never Had Asthma	1,770	26.3	24.0-28.8
No Mammogram within Past Two Years (40+)	282	38.8	32.4-45.6
Mammogram within Past Two Years (40+)	968	19.6	17.0-22.5
No Clinical Breast Exam within Past Two Years	365	33.4	28.1-39.2
Clinical Breast Exam within Past Two Years	871	20.0	17.3-23.1
No Pap Smear within the Past Three Years	255	35.8	29.3-43.0
Pap Smear within the Past Three Years	462	19.5	15.7-23.8
No PSA Test within the Past Two Years (40+)	140	40.3	31.3-50.1
PSA Test within the Past Two Years (40+)	488	23.1	19.2-27.5
No Digital Rectal Exam within the Past Two Years (40+)	179	39.4	31.5-47.9
Digital Rectal Exam within the Past Two Years (40+)	460	22.1	18.1-26.6
Prostate Cancer (40+)	*	*	*
No Prostate Cancer (40+)	565	28.0	24.0-32.3
No Blood Stool Test within the Past Two Years (50+)	1,361	28.1	25.5-31.0
Blood Stool Test within the Past Two Years (50+)	504	19.2	15.6-23.4
Never Had a Sigmoidoscopy or Colonoscopy (50+)	656	37.4	33.2-41.9
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	1,246	19.9	17.5-22.5
Diagnosed with Cancer in the Past 12 Months	*	*	*
Not Diagnosed with Cancer in the Past 12 Months	1,797	25.7	23.5-28.1
Drank Alcohol in Past 30 Days	655	25.2	21.6-29.2
No Alcohol in Past 30 Days	1,291	26.2	23.5-29.0
Binge Drinker	*	*	*
Not a Binge Drinker	1,873	25.4	23.2-27.7
Heavy Drinker	*	*	*
Not a Heavy Drinker	1,882	25.4	23.2-27.7
No Health Insurance (18-64)	-	-	-
Health Insurance (18-64)	-	-	-
Employer Based Health Insurance Coverage (18-64)	_	-	-
Private Health Insurance Plan (18-64)	-	=	-
Medicare (18-64)	-	-	-
Medicaid or Medical Assistance (18-64)	-	-	-
The Military, CHAMPUS, TriCare, or the VA (18-64)	_	-	-
The Indian Health Service (18-64)	-	-	-
No Pneumonia Shot (65+)	648	47.1	42.6-51.6

Table 63 (continued) No Flu Shot, Ages 65 and Older, Within the Past 12 Months for Selected Health Behaviors and Conditions, 2006

Health Behavior or Condition	# Respondents	% No Flu Shot	95% CI
Haven't Been to the Dentist in the Past Year	738	32.5	28.7-36.6
Been to the Dentist in the Past Year	1,200	22.1	19.6-24.9
Not Taking any Precautions Against West Nile Virus	900	30.3	27.0-33.9
Taking Precautions Against West Nile Virus	976	20.9	18.1-23.9
Sometimes/Seldom/Never Wear Seat Belt	257	40.6	33.8-47.8
Always/Almost Always Wear Seat Belt	1,664	23.4	21.2-25.8
Previously Had a Heart Attack	302	24.1	18.9-30.3
Never Had a Heart Attack	1,635	26.3	23.9-28.8
Have Angina or Coronary Heart Disease	260	18.2	13.3-24.3
Do Not have Angina or Coronary Heart Disease	1,654	27.4	25.0-29.9
Previously Had a Stroke	157	23.0	16.4-31.3
Never Had a Stroke	1,790	26.1	23.9-28.5
Fair or Poor Health Status	515	21.1	17.4-25.3
Excellent, Very Good, or Good Health Status	1,425	27.5	24.9-30.3
Physical Health Not Good for 30 Days of the Past 30	211	25.0	19.1-32.1
Physical Health Not Good for 0-29 Days of the Past 30	1,666	25.8	23.5-28.3
Mental Health Not Good for 20-30 Days of the Past 30	*	*	*
Mental Health Not Good for 0-19 Days of the Past 30	1,825	25.4	23.2-27.8
Usual Activities Unattainable for 10-30 Days of the Past 30	150	27.3	19.7-36.4
Usual Activities Unattainable for 0-9 Days of the Past 30	1,757	25.7	23.5-28.1
Dissatisfied / Very Dissatisfied with Life	*	*	*
Satisfied / Very Satisfied with Life	1,836	25.6	23.4-27.9
Physical, Mental, or Emotional Disability	675	22.1	18.7-25.8
No Physical, Mental, or Emotional Disability	1,263	27.9	25.1-30.9
Disability with Special Equipment Needed	329	20.0	15.6-25.2
No Disability with Special Equipment Needed	1,619	27.0	24.6-29.6
Injured in a Fall (45+)	*	*	*
Not Injured in a Fall (45+)	1,820	26.1	23.8-28.5
Never Been Tested for HIV (18-64)	-	-	-
Been Tested for HIV (18-64)	-	-	-
Military Veteran	438	24.7	20.4-29.6
Not a Military Veteran	1,511	26.2	23.8-28.8

Note: *Results based on sample sizes less than 100 have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2006

NO PNEUMONIA SHOT

Definition: Respondents ages 65 and older who have never had a pneumonia vaccination.

Prevalence of No Pneumonia Shot

- South Dakota 35.0%
- o Nationwide median 33.1%


Healthy People 2010 Objective

Decrease the proportion of adults age 65 years and older who have never been vaccinated against pneumococcal disease to 10 percent.

Trend Analysis

Overall, the percent of respondents aged 65 and older who have never had a pneumonia shot has been decreasing. However, in 2006 there was a increase from 33.7 percent in 2005 to 35 percent in 2006. South Dakota has not yet met the *Healthy People 2010 Objective* of 10 percent.

Figure 52 Percent of Respondents, Ages 65 and Older, Who Have Never Had a Pneumonia Shot, 1993-1995, 1997, and 1999-2006



Demographics

Gender There is no significant gender difference in the lack of a pneumonia shot for those

65 and older or for any demographic group between the two.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1993-1995, 1997, and 1999-2006

Age There is a significant decrease in the percentage of those with no pneumonia shot

from the 65-74 age group to the 75 and older age group.

Race There is insufficient data to make any racial comparisons.

Region There are no significant regional differences observed from the available data.

Household Household income does not seem to have a large impact on pneumonia shot **Income** status.

Education Lack of a pneumonia shot seems to generally increase as education increases.

Employment Those who are employed for wages or self-employed demonstrate a very high prevalence of no pneumonia shot, while those who are a homemaker or retired show a very low prevalence.

Marital Those who are married, divorced, or separated exhibit a very high prevalence of Status no pneumonia shot, while those who are widowed show a very low prevalence.

Table 64 Respondents, Ages 65 and Older, Who Have Never Had a Pneumonia Shot, 2006									
Respondents, 2									
	# Resp.	Total %	95% CI	# Resp.	Male %	95% CI	# Resp.	Femal	95% CI
Total	1,889	35.0	(32.6-37.6)	# Kesp.	34.0	(29.9-38.3)	1,257	35.8	(32.8-39.0)
	1,009	33.0	(32.0-37.0)	032	34.0	(29.9-30.3)	1,237	33.0	(32.0-39.0)
<u>Age</u> 18-24									
25-34	-		-	-	-	_		-	-
	-	-	-	-	-	-	-	-	-
35-44	-	-	-	-	-	-	-	-	-
45-54	-	-	-	-	-	-	-	-	-
55-64	- 000	- 10.1	(20.4.46.0)	240	- 42.0	(27.2.40.1)	7.60	- 42.2	(20.5.40.0)
65-74	908	43.1	(39.4-46.9)	340	43.0	(37.2-49.1)	568	43.2	(38.5-48.0)
75+	981	27.8	(24.5-31.2)	292	24.3	(19.2-30.3)	689	29.9	(25.9-34.2)
Race									
White	1,779	34.8	(32.2-37.4)	597	33.8	(29.7-38.2)	1,182	35.4	(32.3-38.7)
American Indian	*	*	*	*	*	*	*	*	*
Region									
Southeast	405	36.9	(32.0-42.1)	127	32.6	(24.8-41.6)	278	39.6	(33.6-46.0)
Northeast	522	34.0	(29.8-38.6)	168	35.0	(27.8-43.0)	354	33.4	(28.3-38.8)
Central	431	31.9	(27.4-36.8)	138	28.9	(21.7-37.5)	293	33.9	(28.4-39.9)
West	387	34.6	(29.7-39.9)	141	37.5	(29.5-46.3)	246	32.0	(26.2-38.5)
American Indian Counties	144	39.3	(30.3-49.1)	*	*	*	*	*	*
Household Income									
Less than \$10.000	115	23.1	(15.4-33.1)	*	*	*	*	*	*
\$10,000-\$14,999	200	33.8	(26.5-42.0)	*	*	*	*	*	*
\$15,000-\$19,999	206	32.4	(25.4-40.3)	*	*	*	*	*	*
\$20,000-\$24,999	231	33.6	(26.9-41.1)	*	*	*	*	*	*
\$25,000-\$34,999	282	40.6	(34.1-47.5)	113	40.2	(30.3-50.9)	169	41.0	(32.7-49.9)
\$35,000-\$49,999	234	37.3	(30.5-44.7)	118	37.4	(28.3-47.6)	116	37.2	(27.6-48.0)
\$50,000-\$74,999	116	25.4	(17.7-35.0)	*	*	*	*	*	*
\$75,000+	*	*	*	*	*	*	*	*	*
Education									
8th Grade or Less	215	33.7	(26.8-41.3)	*	*	*	*	*	*
Some High School	152	39.3	(30.2-49.1)	*	*	*	*	*	*
High School or G.E.D.	731	32.7	(28.8-36.8)	233	33.7	(27.2-40.9)	498	32.0	(27.4-36.9)
	415			233 *	33.7	(27.2-40.9)	490 *	32.0	(27.4-30.9)
Some Post-High School		35.6	(30.5-41.1)					-	
College Graduate	372	38.1	(32.5-44.0)	159	36.9	(29.0-45.6)	213	39.3	(31.8-47.5)
Employment Status	202	7.50	(40.0.54.5)	*	*	*	*	*	*
Employed for Wages	202	56.9	(48.8-64.7)						
Self-employed	176	43.2	(34.9-52.0)	*	*	*	*	*	*
Unemployed	*	*	*	*	*	*	*		
Homemaker	145	39.1	(30.2-48.7)	*	*	*	144	38.5	(29.6-48.2)
Retired	1,297	30.2	(27.4-33.2)	439	27.3	(22.9-32.2)	858	32.4	(28.9-36.2)
Unable to Work	*	*	*	*	*	*	*	*	*
Marital Status									
Married/Unmarried Couple	859	37.0	(33.5-40.6)	425	34.0	(29.3-39.1)	434	40.5	(35.5-45.7)
Divorced/Separated	146	43.0	(33.3-53.2)	*	*	*	*	*	*
Widowed	819	29.3	(26.0-33.0)	115	24.5	(17.0-33.9)	704	30.3	(26.6-34.2)
Never Married	*	*	*	*	*	*	*	*	*


Note: *Results based on sample sizes less than 100 have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2006

National Statistics

The national median for respondents who have never had a pneumonia shot was 33.1 percent. South Dakota had 35.0 percent of respondents who have never had a pneumonia shot. Oregon had the lowest percent of respondents who have never had a pneumonia shot with 25.3 percent, while the District of Columbia had the highest percent with 48 percent.

Figure 53 Nationally, Respondents Ages 65 and Older, Who Have Never Had a Pneumonia Shot, 2006



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2006

Further Analysis

Following are data illustrating the percent of those ages 65 and older who have never had a pneumonia shot for various health behaviors and conditions. For example, 44.0 percent of respondents who stated they sometimes, seldom, or never wear a seat belt had no pneumonia shot, while 33.7 percent of respondents who stated they always or almost always wear a seat belt had no pneumonia shot.

Table 65 No Pneumonia Shot, Ages 65 and Older, for Selected Health Behaviors and Conditions, 2006						
Health Behavior or Condition	# Respondents	% No Pneumonia Shot	95% CI			
Obese (BMI = 30.0+)	398	34.0	28.7-39.8			
Overweight (BMI = 25.0-29.9)	738	34.4	30.6-38.5			
Recommended Weight (BMI = 18.5-24.9)	616	37.2	32.8-41.8			
No Leisure Time Physical Activity	633	35.1	30.8-39.6			
Leisure Time Physical Activity	1,251	35.0	32.0-38.1			
Not Heard of "Healthy South Dakota" Program	1,046	36.2	32.8-39.7			
Heard of "Healthy South Dakota" Program	749	33.8	30.0-37.9			

Table 65 (continued) No Pneumonia Shot, Ages 65 and Older, for Selected Health Behaviors and Conditions, 2006

		% No Pneumonia	
Health Behavior or Condition	# Respondents	Shot	95% CI
Current Smoker	180	43.7	35.3-52.3
Former Smoker	686	29.0	25.2-33.1
Never Smoked	1,021	38.0	34.6-41.6
Smokeless Tobacco Use	*	*	*
No Smokeless Tobacco Use	1,809	34.9	32.3-37.5
Diabetes	288	21.4	16.2-27.7
No Diabetes	1,600	37.4	34.6-40.2
Current Asthma	126	17.4	10.5-27.4
Former Asthma	*	*	*
Never Had Asthma	1,717	36.5	33.9-39.2
No Mammogram within Past Two Years (40+)	275	46.4	39.5-53.4
Mammogram within Past Two Years (40+)	953	32.3	28.9-35.9
No Clinical Breast Exam within Past Two Years	360	42.5	36.6-48.6
Clinical Breast Exam within Past Two Years	855	32.6	29.1-36.4
No Pap Smear within the Past Three Years	253	42.9	36.0-50.1
Pap Smear within the Past Three Years	452	33.2	28.2-38.7
No PSA Test within the Past Two Years (40+)	135	51.7	41.8-61.4
PSA Test within the Past Two Years (40+)	460	28.7	24.3-33.5
No Digital Rectal Exam within the Past Two Years (40+)	170	47.1	38.5-55.9
Digital Rectal Exam within the Past Two Years (40+)	436	28.4	23.9-33.3
Prostate Cancer (40+)	*	*	*
No Prostate Cancer (40+)	533	35.2	30.8-40.0
No Blood Stool Test within the Past Two Years (50+)	1,323	38.8	35.8-41.9
Blood Stool Test within the Past Two Years (50+)	488	23.9	19.7-28.7
Never Had a Sigmoidoscopy or Colonoscopy (50+)	636	50.6	46.0-55.2
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	1,211	27.4	24.6-30.4
Diagnosed with Cancer in the Past 12 Months	*	*	*
Not Diagnosed with Cancer in the Past 12 Months	1,743	35.7	33.1-38.3
Drank Alcohol in Past 30 Days	631	37.7	33.5-42.1
No Alcohol in Past 30 Days	1,254	33.4	30.4-36.6
Binge Drinker	*	*	*
Not a Binge Drinker	1,816	34.8	32.3-37.4
Heavy Drinker	*	*	*
Not a Heavy Drinker	1,822	34.8	32.3-37.4
No Health Insurance (18-64)	-	-	-
Health Insurance (18-64)	-	-	_
Employer Based Health Insurance Coverage (18-64)	-	-	-
Private Health Insurance Plan (18-64)	-	-	-
Medicare (18-64)	-	-	-
Medicaid or Medical Assistance (18-64)	-	-	-
The Military, CHAMPUS, TriCare, or the VA (18-64)	-	-	-
The Indian Health Service (18-64)	-	-	-
No Flu Shot (65+)	509	63.4	58.4-68.0
Flu Shot (65+)	1,378	25.1	22.4-27.9
Haven't Been to the Dentist in the Past Year	717	38.5	34.3-42.8
Been to the Dentist in the Past Year	1,162	33.2	30.1-36.4
Not Taking any Precautions Against West Nile Virus	866	39.0	35.3-42.9
Taking Precautions Against West Nile Virus	953	31.4	28.1-34.9
Sometimes/Seldom/Never Wear Seat Belt	245	44.0	36.8-51.3
Always/Almost Always Wear Seat Belt	1,619	33.7	31.0-36.4

Table 65 (continued) No Pneumonia Shot, Ages 65 and Older, for Selected Health Behaviors and Conditions, 2006

		% No Pneumonia	
Health Behavior or Condition	# Respondents	Shot	95% CI
Previously Had a Heart Attack	293	26.3	20.6-32.9
Never Had a Heart Attack	1,584	36.8	34.0-39.6
Have Angina or Coronary Heart Disease	248	24.6	18.6-31.7
Do Not have Angina or Coronary Heart Disease	1,606	37.3	34.6-40.1
Previously Had a Stroke	151	25.8	18.2-35.3
Never Had a Stroke	1,736	35.8	33.2-38.5
Fair or Poor Health Status	502	24.9	20.7-29.8
Excellent, Very Good, or Good Health Status	1,376	38.5	35.5-41.5
Physical Health Not Good for 30 Days of the Past 30	206	23.9	17.9-31.1
Physical Health Not Good for 0-29 Days of the Past 30	1,611	36.9	34.2-39.7
Mental Health Not Good for 20-30 Days of the Past 30	*	*	*
Mental Health Not Good for 0-19 Days of the Past 30	1,764	34.5	32.0-37.2
Usual Activities Unattainable for 10-30 Days of the Past 30	145	23.7	16.3-33.1
Usual Activities Unattainable for 0-9 Days of the Past 30	1,704	36.2	33.6-38.9
Dissatisfied / Very Dissatisfied with Life	*	*	*
Satisfied / Very Satisfied with Life	1,782	34.9	32.4-37.6
Physical, Mental, or Emotional Disability	659	27.5	23.7-31.7
No Physical, Mental, or Emotional Disability	1,218	39.2	36.0-42.5
Disability with Special Equipment Needed	323	23.4	18.4-29.2
No Disability with Special Equipment Needed	1,564	37.4	34.6-40.2
Injured in a Fall (45+)	*	*	*
Not Injured in a Fall (45+)	1,764	35.6	33.0-38.3
Never Been Tested for HIV (18-64)	-	-	-
Been Tested for HIV (18-64)	-	-	-
Military Veteran	417	29.4	24.7-34.7
Not a Military Veteran	1,471	37.1	34.2-40.0

Note: *Results based on sample sizes less than 100 have been suppressed.

Ssource: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2006